

# The Correlation between Varicocele, Sperm Parameters and Embryo Quality of ICSI Outcome in Infertile Men

Hamid Piroozmanesh<sup>1</sup>, Rahil Jannatifar<sup>2</sup>, Leila Naserpoor<sup>3</sup>, Ebrahim cheraghi<sup>4</sup>

Received 26 September 2020, Accepted for publication 18 January 2021

## Abstract

**Background & Aims:** Varicocele is among the most common identifiable abnormality found in men evaluated for infertility. Despite the long history associated with varicoceles, there remains much controversy regarding their diagnosis and management. The aim of this study was to determine association between varicocele, sperm parameters, and embryo quality of ICSI outcome in infertile men.

**Materials & Methods:** Fifty individuals with varicocele for Intracytoplasmic sperm injection (ICSI) treatment were included in this study. The control group included healthy men without varicoceles (n=50). Semen samples were analyzed according to the World Health Organization (WHO) criteria. Hormonal analysis of serum LH, FSH, and testosterone were measured. Sperm DNA fragmentation was assessed by SCD (Halo sperm), and then ICSI fertilization rate, percentage of high- quality embryos and chemical pregnancy rate were measured.

**Results:** In semen analyses, the total sperm count, total motile sperm and normal sperm morphology were significantly lower in patients with varicocele. Also, the DNA fragmentation was significantly higher in patients with varicocele than patients in the control group ( $p<0.001$ ). The level of reproductive hormones was different in varicocele and control groups ( $p<0.05$ ). Fertilization rate and embryo quality were significantly lower in individuals with varicocele, when compared to men without varicocele ( $p<0.05$ ). The significant negative correlation between DNA fragmentation, fertilization rate, and embryo quality were observed ( $p<0.05$ ).

**Conclusion:** The effect of varicocele on male infertility may be attributed to a decrease in sperm quality as well as an increase in DNA fragmentation, which leads to reduced embryo quality in patients with varicocele.

**Keywords:** Varicocele, DNA fragmentation, Fertilization, Embryo quality

**Address:** Department of Reproductive Biology, the Academic Center for Education, Culture and Research, Qom Branch, Karimi Street, Shahrak Esar, Qom, Iran

**Tel:** +989126526712

**Email:** HP457@yahoo.com

## Introduction

Varicocele, an important factor in the male fertility, can impair sperm quality and fertility(1). Varicocele is the abnormal dilatation of the spermatic veins (2). It is

commonly seen in nearly 40% of male population, affecting 15% of individuals at reproductive age, 35% of those with primary infertility, and up to 80% of men with secondary infertility (3-5).

<sup>1</sup> Department of Reproductive Biology, the Academic Center for Education, Culture and Research, Qom Branch, Qom, Iran (Corresponding Author)

<sup>2</sup> Department of Reproductive Biology, the Academic Center for Education, Culture and Research, Qom Branch, Qom, Iran

<sup>3</sup> Department of Reproductive Biology, the Academic Center for Education, Culture and Research, Qom Branch, Qom, Iran

<sup>4</sup> Department of Biology, Faculty of Science, University of Qom, Qom, Iran

An increase in testicular temperature, scrotal hyperthermia, altered testicular blood flow, testicular hypo perfusion, hypoxia, and testicular hormonal dysfunction may result in varicocele-related testicular dysfunction (6). Recent studies have shown that infertile men with varicocele have abnormal semen parameters (count, motility, and morphology) (7). The correlation between varicocele and dysfunction of spermatogenesis has been well described (8). Perhaps one of the leading causes is high follicle - stimulating hormone (FSH) and low Serum testosterone among infertile men with varicoceles (9, 10).

Varicocele may affect the final stages of spermatogenesis and lead to changes at sperm condensation. During the final stages of spermatogenesis, histones are replaced by cysteine-rich protamines(11). According to one theory, increased thermal caused

damage to the DNA and proteins in the nucleus of spermatid cells and / or Leydig cells (12). Increased testicular temperature as oxidative stress factor is known to have adverse effects on sperm structure and function, such as DNA fragmentation (13). Sperm DNA integrity has been recognized as one of the important determinants of normal fertilization ,embryo development, implantation, and pregnancy rate (14, 15).

ICSI procedure is used to treat severe male factor infertility and to modify their diagnostic and therapeutic approach(16). Also, varicocelectomy have an essential role in the treatment of infertile patients with clinical varicocele and can significantly improve sperm quality and quantity for ART (17, 18). Therefore, we have evaluated the impact of varicoceles on semen quality, reproductive hormones, ICSI outcome and described the associations between DNA fragmentation, fertilization rate, and embryo quality in men with varicocele.

## Method and Material

### Study design and patients:

The ethics committee of Qom University approved the study (IR.QOM.REC.1399.011), and written consent was obtained from the participants. 50 patients who were diagnosed with varicoceles and were referred for infertility treatment at the fertility and infertility of Academic Center for Education, Culture and Research (ACECR), Qom, Iran from October 2018 to December 2019 participated in this study. 50 healthy men without varicoceles were included in the study. Inclusion criteria included male gender, age younger than 40 years, primary infertility, and left-sided varicocele (grades II and III) diagnosed by palpation and Doppler duplex ultrasound. We then extracted physical exam and demographic history from both groups. Semen samples were analyzed according to the World Health Organization (WHO) criteria. Hormonal analysis of serum LH, FSH, and testosterone were measured. Sperm DFI was assessed by Sperm chromatin disruption (SCD) (Halo sperm).

### Sperm collection and semen analysis:

Semen samples were collected by masturbation after 2–5 days of sexual abstinence. After semen liquefaction, sperm parameters analysis was performed according to World Health Organization (19) guidelines, and strict criteria evaluated sperm Morphology after Diff-Quick staining.

### Determination of hormonal analysis:

The hormonal levels in serum of FSH (Human Follicle Stimulating Hormone ELISA Kit (ab108678)(U/ml), LH (Human Follicle Stimulating Hormone (ab108678)(IU/L), and testosterone (Testosterone ELISA Kit (ab108666)(ng/ml) were measured using ELISA (Biotek -ELx 800-enzyme-linked immunosorbent assay).

### Determination of DNA fragmentation (SCD):

For examination of sperm DNA fragmentation, Sperm chromatin dispersion (SCD) test (the Halo sperm

kit, INDAS laboratories, Spain) was used. For each sample, 200 sperms were evaluated under the  $\times 1000$  objective of an optical microscope. In this method, the normal sperm (without fragmented DNA) produce halos (large or medium halos) and abnormal sperm (with fragmented DNA) produce either small halos or no halos (20).

**ICSI technique and embryo culture:**

Oocytes that had the first polar body (MII oocytes) were used for

Intracytoplasmic sperm injection (ICSI). Four hours after oocyte retrieval, a single sperm with normal morphology was injected to inseminate oocyte. At 16–18 hours after ICSI, fertilization was confirmed by the presence of two pronuclei (2PN) under an Olympus inverted microscope (IX71) with a Hoffmann modulation contrast system at  $\times 400$  magnification.

Three-days post- ICSI, embryo quality was assessed based on a three-point scoring system (21): I) symmetric blastomeres and no fragmentation II)

unequal of blastomere’s size and shape and  $<30$  %fragmentation and III) unequal of blastomere’s size and shape and  $>30$  %fragmentation. Chemical pregnancy was defined as serum  $\beta$ hCG  $\geq 20$  IU/L (VIDAS kit) measured 14 days after embryo transfer.

**Statistical Analysis:**

Statistical analysis was performed using SPSS version 20.0 (Chicago, IL, USA). The difference between two independent groups was analyzed using an independent Student’s t - test and mean difference (MD). Significance level was set at  $p < 0.05$

**Results**

**Clinical and demographic characteristics:**

The study population consisted of 50 individuals with grades II and III varicocele. The mean ages of male participants were  $30.1 \pm 4.4$  (range: 22-45) years. There was no statistically significant difference in weight, and Body Mass Index (BMI) between two groups ( $p > 0.05$ ) (Table 1).

**Table 1:** Comparison of the Clinical and demographic characteristics in the varicocele and control groups. ( $p > 0.05$ ).

Clinical and demographic characteristics	Control (n=50)	Varicocele (n=50)	P-value
Age, years	31 (22-35)	32 (22-38)	P= 0.07
Weight, kg	73.0 (59.4–95.0)	73.1 (59.0–93.8)	P=0.11
BMI, kg/m <sup>2</sup>	22.3 (18.7–28.4)	21.4 (18.2–25.8)	P= 0.8
Cigarettes daily	0 (0–20)	0 (0–15)	P= 0.2
Alcohol consumption	0(0–35)	0(0–38)	P=0.10

Data are shown as mean  $\pm$  SE. No difference was observed between the mean of variables in the varicocele group compared with control group. BMI: Body Mass Index.

**Effects of varicocele on seminal parameters:**

Semen analysis revealed that the sperm concentration, total sperm count, total motile sperm were significantly lower in varicocele group than the control group ( $p < 0.001$ ). According to Diff-Quick staining, sperm with normal morphology were significantly higher in the control group, whereas sperm head anomalies were significantly higher in the

varicocele group ( $P < 0.001$ ). When the samples were evaluated with SCD staining, the percentage of DNA fragmentation was  $33 \pm 2.12$  and  $15.1 \pm 1.0$  for the varicocele and control groups, respectively ( $p < 0.001$ ). There was no statistically significant difference in the volume of semen between the groups ( $p > 0.05$ ) (Table 2).

**Table 2:** Comparison of the conventional semen parameters and DNA fragmentation analyses between two groups ( $p<0.05$ ).

Sperm Parameters	Control (n=50)	Varicocele (n=50)	P-value
Volume (ml)	2.6±0.3	2.01±0.1	0.18
Concentration (million/m)	50.5±1.23	28.6±2.13	0.0001
Total sperm count (million/m)	153±12.05	88.4±9.23	0.002
Normal morphology (%)	7.1±0.13	3.5±0.5	0.001
Total Motility (%)	63.23±2.01	39.12±3.2	0.002
DFI (Halo sperm) (%)	15.1±1.03	33±2.12	0.001

Data are shown as mean ± SE. Significant differences for the comparison between two groups are bold type. DFI; DNA Fragmentation Index.

#### Effects of varicocele on fertilization outcome:

The embryo quality was assessed based on a three-point scoring system. I) Cells are of equal size; no fragmentation, II) Cells are of equal size; minor fragmentation only, and III). Cells are of equal or unequal size; fragmentation is moderate to heavy.

As indicated in Table 3, there was a statistically significant difference between the two groups in this

study with regard to the overall fertilization rate and embryo quality ( $p<0.05$ ). Embryo quality includes the number of embryos Grade I ( $p<0.05$ ), number of embryos Grade II ( $p<0.05$ ), number of embryos Grade III ( $p<0.05$ ). There were no statistically significant differences in the clinical pregnancy in varicocele groups compared to control groups ( $p>0.05$ ).

**Table 3:** clinical outcome – comparison between control and varicocele groups ( $p<0.05$ )


Parameters	Control (n=50)	Varicocele (n=50)	P-value
FR (%)	56.9±3.5	34.7±2.2	0.03
No. of embryos Grade I	3.33±0.9	1.53±0.83	0.02
No. of embryos Grade II	3.4±1.2	1.87±0.9	0.02
No. of embryos Grade III	3.8±1.1	1.73±0.8	0.01
No. of clinical pregnancy (%)	33%	28%	0.07

Data are shown as mean ± SE. Significant differences for the comparison between two groups are bold type. FR; Fertilization rate.

#### Effect of varicocele on the level of basal reproductive:

The results of this study showed that testosterone level was significantly lower in the varicocele group

than the control group ( $p<0.05$ ). As well as in this study was showed statistically significantly higher at the levels of FSH, LH hormones in varicocele group than the control group ( $p<0.05$ ).




**Figure 1:** Comparison of (FSH, LH, and testosterone levels) between two groups ( $p < 0.05$ ), \*; Indicate significant difference in varicocele and control groups.

**Correlation between DNA fragmentations, fertilization rate and embryo quality in men with varicocele:**

Data analysis has shown that there is a negative correlation between DNA fragmentation percentage and

fertilization (Fig2A). Sperm DNA fragmentation also was negatively associated with embryo quality in men with varicocele (Fig 2B).



**Figure 2:** Correlation between Fertilization rate (A) and embryo quality (B) with DNA fragmentation index (DFI) in men with varicocele.

## Discussion

Our results demonstrated that patients with the varicocele had lower sperm concentration, total sperm count, and sperm motility compared to the control group, which is consistent with previously reported results (3, 22, 23). The results of this study showed that the number of sperm with abnormal morphology was significantly higher in the varicocele group. The adverse effect of varicocele on sperm quality can be attributed to many factors such as an increased testicular temperature, testicular damage, and hormonal profile abnormalities, and reflux of toxic metabolites from the adrenal glands(24).

One of the critical events in the pathology of varicocele is the excessive production of ROS (25). In terms of pathological conditions, two roles have been envisaged for the overproduction of ROS: ROS-induced damage to the sperm membrane reduces sperm motility and the ability of the sperm to fuse with the oocyte, and ROS directly damages sperm DNA and subsequently affects the genomic integrity of the embryo (26). Oxidative DNA damage caused by the extent of oxidative stress (OS) led to a reduction in sperm quality in patients with varicoceles (27).

According to the results, the data has demonstrated that there is an increase in DNA fragmentation in the sperm of the varicocele group compared to the control group. In this regard, several pieces of evidence have demonstrated the relationship between varicocele and sperm DNA damage (28, 29). The final steps of spermatogenesis have an important role in sperm function, and fertility. Disorders affecting varicocele on sperm DNA denaturation in the later stages of spermatogenesis may lead to abortive apoptosis or altered fertility potential (30).

There are many aspects still unknown today about the effects of varicocele on hormonal levels (31). Higher levels of FSH and LH indicate that a subtle Leydig cell

dysfunction was associated with varicocele and may cause changes in Sertoli cellular function, and decrease testosterone production by Leydig cells (32, 33). Following other authors (10, 34), our study shows that men with varicocele had a higher LH and FSH concentration, although testosterone level was significantly lower than the control group. These differences in male reproductive hormones seen in infertile patients with varicoceles could be related to the lower sperm concentration and lower sperm motility (35). In our study, we found that in patients with varicocele ICSI resulted (fertilization rate and embryo development) significantly lower compared to fertile men. Therefore, varicocele might have a negative impact on fertilization rate and embryo quality following ICSI. Also, a negative correlation with sperm DNA fragmentation, fertilization rate, and embryo quality is observed in varicocele patients. The negative effect of sperm DNA damage on embryo development might have been modulated by the ability of the oocyte to repair sperm DNA damage before the first cleavage (36, 37). Recent studies evaluated that abnormal sperm chromatin packaging is correlated with the reduced ability of spermatozoa to fertilize oocytes in standard, conception or during ART procedures (38-40). A high percentage of DNA fragmentation index values in varicocele patients may be associated with a lower fertilization and embryo quality in assisted reproductive techniques.

The conclusions outlined above mainly support the varicocele repair (Varicocelectomy), and antioxidant therapy may benefit sperm quality and success for couples with varicocele-related infertility who require ART to initiate a pregnancy (41). Barekat et al. (42) suggested the use of N-acetyl-L-cysteine, as an antioxidant, after varicocelectomy, and reported that post-operation antioxidant treatment reduced ROS levels and improved chromatin integrity and pregnancy

rates. Therefore, varicocele repair should be offered as part of treatment for infertile couples with palpable varicoceles.

### Conclusion

In conclusion, Varicocele remains a topic with many controversies for the couples seeking fertility. Our clinical study of infertile men showed that semen quality was significantly impaired in men with varicocele. Despite a decrease in fertilization rate and embryo quality transfer in varicocele patients, no significant effect on clinical pregnancy was observed.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgements

This study was supported by IVF Unit of Infertility Research Center of the ACECR, Qom.

### References

1. Esteves SC, Miyaoka R, Agarwal AJC. An update on the clinical assessment of the infertile male. *Clinics* 2011;66(4):691-700.
2. Shiraishi K, Matsuyama H, Takihara HJJjou. Pathophysiology of varicocele in male infertility in the era of assisted reproductive technology. *Int J Urol* 2012;19(6):538-50.
3. Jensen CFS, Østergren P, Dupree JM, Ohl DA, Sønksen J, Fode MJNRU. Varicocele and male infertility. *Nature Reviews Urology* 2017;14(9):523.
4. Rotker K, Sigman MJAjoa. Recurrent varicocele. *Asian journal of andrology* 2016;18(2):229.
5. Jarow JP, Coburn M, Sigman MJU. Incidence of varicoceles in men with primary and secondary infertility. *Urology* 1996;47(1):73-6.
6. Alsaikhan B, Alrabeeah K, Delouya G, Zini AJAjoa. Epidemiology of varicocele. *Asian journal of andrology* 2016;18(2):179.
7. Pasqualotto FF, Lucon AM, de Góes PM, Sobreiro BP, Hallak J, Pasqualotto EB, et al. Semen profile, testicular volume, and hormonal levels in infertile patients with varicoceles compared with fertile men with and without varicoceles. *Fertil Steril* 2005;83(1):74-7.
8. Fuse H, Iwasaki M, Mizuno I, Ikehara-Kawauchi YJAoa. Evaluation of acrosome reactivity using the Acrobeads test in varicocele patients: findings before and after treatment. *Archives of andrology* 2003;49(1):1-6.
9. Kumanov P, Nandipati K, Tomova A, Agarwal AJF. Inhibin B is a better marker of spermatogenesis than other hormones in the evaluation of male factor infertility. *Fertil Steril* 2006;86(2):332-8.
10. Damsgaard J, Joensen UN, Carlsen E, Erenpreiss J, Jensen MB, Matulevicius V, et al. Varicocele is associated with impaired semen quality and reproductive hormone levels: a study of 7035 healthy young men from six European countries. *Eur Urol* 2016;70(6):1019-29.
11. Roque M, Esteves SC. Effect of varicocele repair on sperm DNA fragmentation: a review. *Int Urol Nephrol* 2018;50(4):583-603.
12. Naughton CK, Nangia AK, Agarwal AJHru. Varicocele and male infertility: part II: pathophysiology of varicoceles in male infertility. *Hum Reprod* 2001;7(5):473-81.
13. Bisht S, Dada RJFB. Oxidative stress: Major executioner in disease pathology, role in sperm DNA damage and preventive strategies. *Front Biosci (Schol Ed)* 2017;9:420-47.
14. Kim SM, Kim SK, Jee BC, Kim SH. Effect of sperm DNA fragmentation on embryo quality in normal responder women in in vitro fertilization and intracytoplasmic sperm injection. *Yonsei medical journal* 2019;60(5):461-6.
15. Zheng W-W, Song G, Wang Q-L, Liu S-W, Zhu X-L, Deng S-M, et al. Sperm DNA damage has a negative effect on early embryonic development following in vitro fertilization. *Asian journal of andrology* 2018;20(1):75.

16. Kupka MS, Dorn C, Richter O, Felberbaum R, van der Ven HJF. Impact of reproductive history on in vitro fertilization and intracytoplasmic sperm injection outcome: evidence from the German IVF Registry. *Fertil Steril* 2003;80(3):508-16.
17. Gabrielsen JS, Thirumavalavan N, Pastuszak AWJV, Guide MIAC. Effect of Varicocele Treatment. 2019:295.
18. Pathak P, Chandrashekar A, Hakky TS, Pastuszak AWJAjoa. Varicocele management in the era of in vitro fertilization/intracytoplasmic sperm injection. *Asian journal of andrology* 2016;18(3):343.
19. Organization WH. World health statistics 2010: World Health Organization; 2010.
20. Yilmaz S, Demiroglu ZA, Yilmaz E, Sofuoglu K, Delikara N, Kutlu P. Effects of sperm DNA fragmentation on semen parameters and ICSI outcome determined by an improved SCD test, Halosperm. *International Journal of Fertility and Sterility* 2010; 4(2): 73-8.
21. Ding J, Xu T, Tan X, Jin H, Shao J, Li HJE, et al. Raman spectrum: A potential biomarker for embryo assessment during in vitro fertilization. *Exp Ther Med* 2017;13(5):1789-92.
22. Hauser R, Paz G, Botchan A, Yogev L, Yavetz H. Varicocele and male infertility: part II: varicocele: effect on sperm functions. *Human Reproduction Update* 2001;7(5):482-5.
23. Baazeem A, Belzile E, Ciampi A, Dohle G, Jarvi K, Salonia A, et al. Varicocele and male factor infertility treatment: a new meta-analysis and review of the role of varicocele repair. *Eur Urol* 2011;60(4):796-808.
24. Kantartzi P, Goulis CD, Goulis G, Papadimas I. Male infertility and varicocele: myths and reality. *Hippokratia* 2007;11(3):99.
25. Agarwal A, Prabakaran S, Allamaneni SS. Relationship between oxidative stress, varicocele and infertility: a meta-analysis. *Reprod Biomed Online* 2006;12(5):630-3.
26. Cho C-L, Esteves SC, Agarwal A. Novel insights into the pathophysiology of varicocele and its association with reactive oxygen species and sperm DNA fragmentation. *Asian Journal of Andrology* 2016;18(2):186.
27. Kohn JR, Haney NM, Nichols PE, Rodriguez KM, Kohn TP. Varicocele Repair Prior to Assist Reproductive Technology: Patient Selection and Special Considerations. *Res Rep Urol* 2020;12:149.
28. Wang Y-J, Zhang R-Q, Lin Y-J, Zhang R-G, Zhang W-L. Relationship between varicocele and sperm DNA damage and the effect of varicocele repair: a meta-analysis. *Reprod Biomed Online* 2012;25(3):307-14.
29. Esfahani MHN, Tavalae M. Origin and role of DNA damage in varicocele. *International journal of fertility & sterility* 2012;6(3):141.
30. Zümürütbaş AE, Gülpınar Ö, Mermerkaya M, Süer E, Yaman Ö. The effect of varicocele on sperm morphology and DNA maturity: does acridine orange staining facilitate diagnosis? *Turk J Urol* 2013;39(3):165.
31. Tanrikut C, Goldstein M. Varicocele repair for treatment of androgen deficiency. *Current opinion in urology* 2010;20(6):500-2.
32. Luo D-Y, Yang G, Liu J-J, Yang Y-R, Dong Q. Effects of varicocele on testosterone, apoptosis and expression of StAR mRNA in rat Leydig cells. *Asian journal of andrology* 2011;13(2):287.
33. Dabaja A, Wosnitzer M, Goldstein M. Varicocele and hypogonadism. *Current urology reports* 2013;14(4):309-14.
34. Çayan S, Akbay E, Saylam B, Kadioğlu A. Effect of varicocele and its treatment on testosterone in hypogonadal men with varicocele: review of the literature. *Balkan medical journal* 2020;37(3):121.
35. Niederberger CJTJou. Re: Effects of Varicocele on Serum Testosterone and Changes of Testosterone after Varicolectomy: A Prospective Controlled Study. 2015.
36. Simon L, Murphy K, Shamsi M, Liu L, Emery B, Aston K, et al. Paternal influence of sperm DNA integrity on early embryonic development. *Hum Reprod* 2014;29(11):2402-12.



37. Meseguer M, Martinez-Conejero J, O'Connor JE, Pellicer A, Remohí J, Garrido NJF, et al. The significance of sperm DNA oxidation in embryo development and reproductive outcome in an oocyte donation program: a new model to study a male infertility prognostic factor. *Fertil Steril* 2008;89(5):1191-9.
38. Sönmez MG, Haliloğlu AH. Role of varicocele treatment in assisted reproductive technologies. *Arab journal of urology* 2018;16(1):188-96.
39. Kohn TP, Kohn JR, Pastuszak AW. Varicolectomy before assisted reproductive technology: are outcomes improved? *Fertil Steril* 2017;108(3):385-91.
40. Coward RM. Evolving role of varicocele repair in the era of assisted reproduction. *Fertil Steril* 2017;108(4):596-7.
41. Steiner AZ, Hansen KR, Barnhart KT, Cedars MI, Legro RS, Diamond MP, et al. The effect of antioxidants on male factor infertility: the Males, Antioxidants, and Infertility (MOXI) randomized clinical trial. *Fertil Steril* 2020 ;113(3):552-60.
42. Barekat F, Tavalae M, Deemeh MR, Bahreinian M, Azadi L, Abbasi H, et al. A preliminary study: N-acetyl-L-cysteine improves semen quality following varicolectomy. *International journal of fertility & sterility* 2016;10(1):120.